

April, 1992
BIR/PH/92-2.

CLASSICAL HAMILTONIAN REDUCTION AND SUPERCONFORMAL ALGEBRAS

W. A. SABRA[★]

*Physics Department
Birkbeck College,
Malet Street
London WC1E 7HX*

ABSTRACT

The Polyakov's "soldering procedure" which shows how two-dimensional diffeomorphisms can be obtained from $SL(2, \mathbb{R})$ gauge transformations is discussed using the free-field representation of $SL(2, \mathbb{R})$ current algebra. Using this formalism, the relation of Polyakov's method to that of the Hamiltonian reduction becomes transparent. This discussion is then generalised to $N=1$ superdiffeomorphisms which can be obtained from $N=1$ super $Osp(1,2)$ gauge transformations. It is also demonstrated that the phase space of the $Osp(2,2)$ supercurrent algebra represented by free superfields is connected to the classical phase space of $N=2$ superconformal algebra via Hamiltonian reduction.

★ email ubap734@uk.ac.bbk.cu

1. Introduction

In recent years, the Hamiltonian reduction of the phase space of the Wess-Zumino-Novikov-Witten (WZNW) models with the gauge group $SL(2, \mathbb{R})$ and its graded extensions [1-6] provided more understanding not only of the structure of conformal and superconformal algebras but also of the origin of non-compact current algebra symmetries in the chiral gauge formulation of induced two dimensional gravity and supergravity theories [7]. Classically, the Hamiltonian reduction [3, 4] can be described as follows. A constraint is imposed on the affine currents describing the phase space of the WZNW model. The constrained phase space modulo gauge transformations, generated by the residual symmetry preserving the constraints, is the phase space of the reduced theory.

In [4] by imposing a certain constraint on the $SL(2, \mathbb{R})$ Kac-Moody current algebra, an irreducible representation of the Virasoro algebra was obtained from an irreducible representation of the $SL(2, \mathbb{R})$ Kac-Moody current algebra using the BRST formalism. This method is the quantum analogue of the Hamiltonian reduction. The hidden $SL(2, \mathbb{R})$ current algebra symmetry in the Virasoro algebra manifests itself in the chiral gauge formulation of induced two dimensional quantum gravity [7]. The results of [4] were extended to the supersymmetric cases in [5], in which the $N=1$ and $N=2$ superconformal algebras were obtained as the Hamiltonian reduction of the bosonic Kac-Moody current algebras of $Osp(1,2)$ and $Osp(2,2)$ respectively. In this construction, extra fermionic degrees of freedom have to be introduced in order to get a consistent algebra of constraints.

In refs. [8,9], following the method of Polyakov in which the conformal transformation of the stress energy tensor defining the phase space of Virasoro algebra was derived from the restricted $SL(2, \mathbb{R})$ gauge transformations, the $N=1$ and $N=2$ superconformal transformations were derived as restricted $N=1$ $Osp(1,2)$ and $Osp(2,2)$ supergauge transformations respectively. The method of Polyakov is simply a Hamiltonian reduction in disguise. This is best explained by using free fields (Wakimoto fields) [10,11] to describe the phase space of $SL(2, \mathbb{R})$ WZNW model.

Introduce a scalar field ϕ and a pair of bosonic ghosts (β, γ) of dimensions $(1, 0)$. The Poisson brackets of these fields are given by^{*}

$$\begin{aligned}\{\beta(z_1), \gamma(z_2)\}_{P.B} &= -\delta(z_1 - z_2), \\ \{\partial_{z_1}\phi(z_1), \phi(z_2)\}_{P.B} &= -\frac{1}{2k}\delta(z_1 - z_2).\end{aligned}\tag{1.1}$$

The $SL(2, R)$ conserved classical currents can be written as

$$\begin{aligned}J^+ &= k\partial_z\gamma + 2k\gamma\partial_z\phi - \gamma^2\beta, \\ J^- &= \beta, \\ J^0 &= -\beta\gamma + k\partial_z\phi,\end{aligned}\tag{1.2}$$

where k is the level of the $SL(2, R)$ current algebra and $\{0, +, -\}$ are the indices of the $SL(2, R)$ group. One way to construct the currents in terms of the free fields is to set $J^- = \beta$ and write the most general form of J^0 and J^+ in terms of the free fields β, γ and $\partial_z\phi$, which is then fixed by demanding that the Poisson brackets of the currents satisfy the classical $SL(2, R)$ current algebra [11]. The currents can also be derived from the $SL(2, R)$ WZNW model as follows. One can parametrize the group element g of $SL(2, R)$ by the Gauss product [2]

$$g = \begin{pmatrix} 1 & 0 \\ \gamma & 1 \end{pmatrix} \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix} \begin{pmatrix} 1 & F \\ 0 & 1 \end{pmatrix}.\tag{1.3}$$

Then the $SL(2, R)$ WZNW action [12,13,14] can be written as

$$S \sim k \int d^2z \left(\frac{1}{\lambda^2} \partial_z \lambda \partial_{\bar{z}} \lambda + \lambda^2 \partial_z F \partial_{\bar{z}} \gamma \right).\tag{1.4}$$

The right-moving conserved currents derived from the action (1.4) are given by

$$\begin{aligned}J &= \begin{pmatrix} J^0 & J^- \\ J^+ & -J^0 \end{pmatrix} = k\partial_z g g^{-1} \\ &= k \begin{pmatrix} \partial_z \lambda / \lambda - \gamma \lambda^2 \partial_z F & \lambda^2 \partial_z F \\ \partial_z \gamma + 2\gamma \partial_z \lambda / \lambda - \gamma^2 \lambda^2 \partial_z F & -\partial_z \lambda / \lambda + \gamma \lambda^2 \partial_z F \end{pmatrix}.\end{aligned}\tag{1.5}$$

^{*} we suppress the space-time indices of the fields and parametrize the two dimensional space time with coordinates (x, t) by $z = t + x$ and $\bar{z} = t - x$.

By setting

$$k\lambda^2\partial_z F = \beta, \quad \lambda = e^\phi, \quad (1.6)$$

we get the expressions of the currents in (1.2) and the action (1.4) becomes an action of a free scalar field and a bosonic ghost system,

$$S \sim \int d^2z \left(k\partial_z\phi\partial_{\bar{z}}\phi + \beta\partial_{\bar{z}}\gamma \right). \quad (1.7)$$

Polyakov's partial gauge fixing [6] is given as

$$J^- = k, \quad J^0 = 0, \quad J^+ = -T. \quad (1.8)$$

Solving (1.8) in terms of the free fields gives

$$T = -k(\partial_z^2\phi + (\partial_z\phi)^2), \quad (1.9)$$

which is recognized as the classical Feign-Fuchs equation [16] for the spin-2 generator describing the phase space of Virasoro algebra. The Poisson bracket of T is calculated using (1.1) and is given by

$$\{T(z_1), T(z_2)\}_{\text{P.B.}} = \frac{k}{2}\partial_{z_2}^3\delta(z_1 - z_2) + 2T(z_2)\partial_{z_2}\delta(z_1 - z_2) + \partial_{z_2}T(z_2)\delta(z_1 - z_2). \quad (1.10)$$

The reason why J^+ becomes a spin-2 field was explained in [6] as a result of the “soldering” of the isospin space and the two dimensional space time in the background $J^- = k$. In terms of the free fields this can be explained as a result of the fact that after the partial gauge fixing, the field γ originally of spin-0 becomes a spin-1 field ($\gamma = \partial_z\phi$). In the classical Hamiltonian reduction method [2,4], the constraint $J^- = k$ is first imposed. Then the constrained phase space has a one-parameter residual symmetry generated by the Borel subgroup of $\text{SL}(2, \mathbb{R})$, this residual symmetry can be used to gauge away a further degree of freedom giving a

reduced theory with one degree of freedom. An element of the space of the residual symmetry that preserves the constraint $J^- = k$ is given by

$$u = \begin{pmatrix} 1 & 0 \\ a & 1 \end{pmatrix}, \quad (1.11)$$

where a is a gauge parameter. Under the finite residual gauge transformation

$$g \rightarrow ug, \quad (1.12)$$

The currents transform as

$$J \rightarrow uJu^{-1} + k\partial_z uu^{-1}. \quad (1.13)$$

Exploiting the gauge symmetry (1.13) and selecting the gauge parameter as (Drinfeld-Sokolov gauge)

$$a = \frac{1}{k}J^0, \quad (1.14)$$

one can put J into the form

$$J_{DS} = \begin{pmatrix} 0 & k \\ -T & 0 \end{pmatrix}, \quad (1.15)$$

where T , the coordinate of the reduced phase space is given by Eq.(1.9). This demonstrates that Polyakov's partial gauge fixing is equivalent to the Drinfeld-Sokolov gauge in the Hamiltonian reduction method.

It has to be mentioned that there also exists another gauge choice, the so called diagonal gauge in which the reduced space takes the form

$$\mathbf{J}_{diagonal} = k \begin{pmatrix} \partial_z \phi & 1 \\ 0 & -\partial_z \phi \end{pmatrix}. \quad (1.16)$$

The connection between the coordinates of the reduced space in the two different gauges can be explained as follows. Consider the system of linear differential

equations

$$(k\partial_z - J) \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = 0, \quad (1.17)$$

where v_1 and v_2 are the components of a two-dimensional vector. In the above system of differential equations one can eliminate the component v_2 and obtain a gauge invariant second order differential equation for v_1 . Computing (1.17) in the diagonal gauge gives

$$\begin{aligned} k\partial_z v_1 - k\partial_z \phi v_1 - kv_2 &= 0, \\ k\partial_z v_2 + k\partial_z \phi v_2 &= 0, \end{aligned} \quad (1.18)$$

after eliminating v_2 in (1.18) we get

$$k\partial_z^2 v_1 - k\partial_z \phi \partial_z \phi v_1 - k\partial_z^2 \phi v_1 = 0 \quad (1.19)$$

The reduced space of Drinfeld-Sokolov gauge can be deduced from that of the diagonal gauge as follows. Assuming we put the reduced space to the form

$$J = \begin{pmatrix} 0 & k \\ -T & 0 \end{pmatrix}. \quad (1.20)$$

Then computing (1.17) using this form of J , we obtain (after solving for v_2 in terms of v_1) the differential equation

$$k\partial_z^2 v_1 + Tv_1 = 0. \quad (1.21)$$

By identifying equations (1.19) and (1.21), we arrive at the form of T . The relation which connects T to the free field $\partial_z \phi$ is known as the Miura transformation [3,4].

The analysis of [8] suggests that the phase space of of N=1 Osp(1,2) current algebras can be reduced to that of N=1 superconformal algebra. A similar observation has been made in [15] in which a quantum Hamiltonian reduction of super Osp(1,2) Kac-Moody algebra has been performed.

This work is organized as follows. In the next section, the results of [8] in which Polyakov's "soldering procedure" is used to show how two-dimensional N=1 superdiffeomorphisms can be obtained from N=1 Osp(1,2) gauge transformations is discussed using the free-superfield realization of N=1 Osp(1,2) supercurrent algebra. In section 3, we construct the free-superfield realization of N=1 Osp(2,2) supercurrent algebra. Using this representation, we impose a certain set of constraints on the phase space of the supercurrent algebra and perform the classical Hamiltonian reduction of the system. The resulting phase space is shown to have the N=2 superconformal symmetry. Section 4 contains a summary of our results and a discussion of the quantum theory.

2. Classical N=1 superVirasoro algebra

In this section the free-superfield representation of the Osp(1,2) supercurrents [5] is used to verify the results of [8] in which the N=1 superconformal transformations were derived as restricted N=1 Osp(1,2) supergauge transformations. The N=1 Osp(1,2) supercurrent algebra can be described in terms of one scalar superfield Φ , two pairs of superghosts (B, C) and $(\Psi^\dagger, \Psi)^*$ of dimensions $(1/2, 0)$ as

$$\begin{aligned}
\mathbf{J}^+ &= kDC + 2kCD\Phi + 2C\Psi\Psi^\dagger + C^2B - kD\Psi\Psi, \\
\mathbf{J}^- &= -B, \\
\mathbf{J}^0 &= BC + kD\Phi + \Psi\Psi^\dagger, \\
\mathbf{J}^{-\frac{1}{2}} &= \frac{1}{2}B\Psi - \Psi^\dagger, \\
\mathbf{J}^{+\frac{1}{2}} &= kD\Psi - k\Psi D\Phi + C\mathbf{J}^{-\frac{1}{2}},
\end{aligned} \tag{2.1}$$

where $D = \partial_\theta + \theta\partial_z$ and $\{0, +, -, -\frac{1}{2}, +\frac{1}{2}\}$ are the Osp(1,2) group indices. Imposing

* The superghosts in components are expressed as $B = b - \theta\beta$, $C = \gamma + \theta c$ where (b, c) are fermionic ghosts and (β, γ) are bosonic. Similarly $\Psi^\dagger = \alpha + \theta\psi^\dagger$ and $\Psi = \psi + \theta\delta$ where the ghosts $(\psi^\dagger, \psi)$ are fermionic and (α, δ) are bosonic. We mention that it is the ghosts (β, γ) and $(\psi^\dagger, \psi)$ that appear in the free-field realization of the bosonic Osp(1,2) current algebra [5].

the partial gauge fixing of [8]

$$\mathbf{J}^- = \mathbf{J}^0 = \mathbf{J}^{+\frac{1}{2}} = 0, \quad \mathbf{J}^{-\frac{1}{2}} = k, \quad \mathbf{J}^+ = -\mathbf{T}, \quad (2.2)$$

gives the following relations among the free superfields

$$B = 0, \quad \Psi^\dagger = -k, \quad D\Phi = \Psi, \quad C = -\partial_z \Phi. \quad (2.3)$$

These relations when substituted back in $\mathbf{T}$ give,

$$\mathbf{T} = k(\partial_z \Phi D\Phi + \partial_z D\Phi). \quad (2.4)$$

Obviously $\mathbf{T}$ is the Feign-Fuchs representation of a spin-3/2 superfield describing the phase space of N=1 superVirasoro algebra. This can be easily demonstrated by computing the Poisson brackets of the field components of $\mathbf{T}$. Writing

$$\mathbf{T} = G - \theta T, \quad \Phi = \phi + \theta\psi, \quad (2.5)$$

and using

$$\{D\Phi(z_1, \theta_1), \Phi(z_2, \theta_2)\}_{\text{P.B}} = -\frac{1}{2k}\theta_{12}\delta(z_{12}), \quad (2.6)$$

where

$$\theta_{12} = \theta_1 - \theta_2, \quad z_{12} = z_1 - z_2 - \theta_1\theta_2, \quad (2.7)$$

the Poisson brackets of the component fields

$$T = -k(\partial_z^2 \phi + \partial_z \psi\psi + (\partial_z \phi)^2), \quad G = k(\partial_z \phi\psi + \partial_z \psi) \quad (2.8)$$

can be shown to satisfy the classical N=1 superVirasoro algebra,

$$\begin{aligned} \{T(z_1), T(z_2)\}_{\text{P.B}} &= \frac{k}{2}\partial_{z_2}^3 \delta(z_1 - z_2) + 2T(z_2)\partial_{z_2} \delta(z_1 - z_2) + \partial_{z_2} T(z_2)\delta(z_1 - z_2), \\ \{T(z_1), G(z_2)\}_{\text{P.B}} &= \delta(z_1 - z_2)\partial_{z_2} G(z_2) + \frac{3}{2}\partial_{z_2} \delta(z_1 - z_2)G(z_2), \\ \{G(z_1), G(z_2)\}_{\text{P.B}} &= \frac{k}{2}\partial_{z_2}^2 \delta(z_1 - z_2) + \frac{1}{2}\delta(z_1 - z_2)T(z_2). \end{aligned} \quad (2.9)$$

In the Hamiltonian reduction method, the partial gauge fixing (2.2) is equivalent

to the Drinfeld-Sokolov gauge, i.e, imposing the constraints

$$\mathbf{J}^- = 0, \quad \mathbf{J}^{-\frac{1}{2}} = k \quad (2.10)$$

and then using the residual gauge transformations (generated by the Borel super subgroup) to set $\mathbf{J}^0$ and $\mathbf{J}^{+\frac{1}{2}}$ to zero. However, in the diagonal gauge one uses the residual gauge transformations to set $\mathbf{J}^+$ and $\mathbf{J}^{+\frac{1}{2}}$ to zero. In this case the reduced space is described by the free coordinate $D\Phi$. The coordinates $\mathbf{T}$ and $D\Phi$ are connected by the super Miura transformation. This connection can be explained as follows. Consider the system of linear differential equation

$$kDv - \begin{pmatrix} \mathbf{J}^0 & 0 & k \\ \mathbf{J}^+ & -\mathbf{J}^0 & \mathbf{J}^{+\frac{1}{2}} \\ \mathbf{J}^{+\frac{1}{2}} & -k & 0 \end{pmatrix} v = 0, \quad v = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} \quad (2.11)$$

where v is a vector. This system of differential equations can be written into a higher order gauge-invariant differential equation in terms of the component v_1 . Computing (2.11) in the diagonal gauge gives

$$\begin{aligned} kDv_1 - kD\Phi v_1 - kv_3 &= 0, \\ kDv_2 + kD\Phi v_2 &= 0, \\ kDv_3 + kv_2 &= 0. \end{aligned} \quad (2.12)$$

By solving these equation in terms of v_1 we obtain

$$k\partial_z Dv_1 - k(\partial_z D\Phi + \partial_z \Phi D\Phi)v_1 = 0. \quad (2.13)$$

In the Drinfeld-Sokolov gauge one obtains from (2.11) the differential equations

$$\begin{aligned} kDv_1 - kv_3 &= 0, \\ kDv_2 + \mathbf{T}v_1 &= 0, \\ kDv_3 + kv_2 &= 0. \end{aligned} \quad (2.14)$$

Solving these equations in terms of v_1 we get

$$k\partial_z Dv_1 - \mathbf{T}v_1 = 0. \quad (2.15)$$

Comparing (2.13) and (2.15) we get the expression of $\mathbf{T}$. For a discussion of the quantum hamiltonian reduction of N=1 Osp(1,2) supercurrent algebra the reader is referred to [15].

3. Classical N=2 superVirasoro algebra

In this section, motivated by the construction of [9] in which it is shown that N=2 superdiffeomorphisms are restricted N=1 supergauge transformations of a supergauge theory with Osp(2,2) as a gauge group, we will consider in some details the Hamiltonian reduction of the N=1 Osp(2,2) supercurrent algebra using its free-superfield realization.

3.1. FREE- SUPERFIELD REALIZATION OF N=1 OSP(2,2) SUPERCURRENT ALGEBRA

In this section an explicit construction of the free superfields describing the phase space of N=1 Osp(2,2) WZNW model is given. The orthosymplectic group Osp(2,2) is generated by four bosonic generators $\{l_0, l_{-1}, l_1, l_u\}$ and four fermionic generators $\{(l_{\frac{1}{2}})_1, (l_{\frac{1}{2}})_2, (l_{-\frac{1}{2}})_1, (l_{-\frac{1}{2}})_2\}$, which are represented as follows,

$$l_0 = \begin{pmatrix} 1/2 & 0 & 0 & 0 \\ 0 & -1/2 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad l_1 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad l_{-1} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$l_u = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix} \quad (l_{\frac{1}{2}})_1 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (l_{\frac{1}{2}})_2 = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}$$

$$(l_{-\frac{1}{2}})_1 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (l_{-\frac{1}{2}})_2 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}. \quad (3.1)$$

A group element $g \in \text{Osp}(2,2)$ can be represented by

$$g = \begin{pmatrix} 1 & 0 & 0 & 0 \\ C & 1 & \Psi_1 & \Psi_2 \\ \Psi_1 & 0 & 1 & 0 \\ \Psi_2 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \lambda & 0 & 0 & 0 \\ 0 & \lambda^{-1} & 0 & 0 \\ 0 & 0 & G_1 & G_2 \\ 0 & 0 & -G_2 & G_1 \end{pmatrix} \begin{pmatrix} 1 & F & \xi_1 & \xi_2 \\ 0 & 1 & 0 & 0 \\ 0 & -\xi_1 & 1 & 0 \\ 0 & -\xi_2 & 0 & 1 \end{pmatrix}, \quad (3.2)$$

where $\{C, \lambda, F, G_1, G_2\}$ are bosonic superfields ($G_1^2 + G_2^2 = 1$) and $\{\Psi_1, \Psi_2, \xi_1, \xi_2\}$ are fermionic superfields. The N=1 super $\text{Osp}(2,2)$ WZNW action [17,18] can then be expressed as

$$\begin{aligned} S \sim k \int d^2 z d^2 \theta & \left(\bar{D}\Phi_0 D\Phi_0 + \bar{D}\Phi_1 D\Phi_1 \right. \\ & + \lambda^2 (\bar{D}C - \Psi_1 \bar{D}\Psi_1 - \Psi_2 \bar{D}\Psi_2) (DF + D\xi_1 \xi_1 + D\xi_2 \xi_2) \\ & \left. - 2\lambda \bar{D}\Psi_1 (G_1 D\xi_1 + G_2 D\xi_2) - 2\lambda \bar{D}\Psi_2 (G_1 D\xi_2 - G_2 D\xi_1), \right) \end{aligned} \quad (3.3)$$

where the covariant derivatives D and $\bar{D}$ are given in terms of the holomorphic and anti-holomorphic Grassmann coordinates θ and $\bar{\theta}$ by

$$D \equiv \frac{\partial}{\partial \theta} + \theta \frac{\partial}{\partial z}, \quad \bar{D} \equiv \frac{\partial}{\partial \bar{\theta}} + \bar{\theta} \frac{\partial}{\partial \bar{z}}. \quad (3.4)$$

The holomorphic supercurrents of the super WZNW model are given by

$$\mathbf{J} = k D g g^{-1} = \begin{pmatrix} \mathbf{J}^0 & \mathbf{J}^- & \mathbf{J}_1^{-\frac{1}{2}} & \mathbf{J}_2^{-\frac{1}{2}} \\ \mathbf{J}^+ & -\mathbf{J}^0 & \mathbf{J}_1^{+\frac{1}{2}} & \mathbf{J}_2^{+\frac{1}{2}} \\ \mathbf{J}_1^{+\frac{1}{2}} & -\mathbf{J}_1^{-\frac{1}{2}} & 0 & -\mathbf{J}^u \\ \mathbf{J}_2^{+\frac{1}{2}} & -\mathbf{J}_2^{-\frac{1}{2}} & \mathbf{J}^u & 0 \end{pmatrix}. \quad (3.5)$$

If the following change of superfields is performed

$$\begin{aligned}
B &= -k\lambda^2(DF + D\xi_1\xi_1 + D\xi_2\xi_2), \\
\Psi_1^\dagger &= \frac{1}{2}B\Psi_1 - k\lambda(G_1D\xi_1 + G_2D\xi_2), \\
\Psi_2^\dagger &= \frac{1}{2}B\Psi_2 - k\lambda(G_1D\xi_2 - G_2D\xi_1),
\end{aligned} \tag{3.6}$$

then the super WZNW action in (3.3) becomes an action of free superfields

$$S \sim \int d^2z d^2\theta \left(k\bar{D}\Phi_0 D\Phi_0 + k\bar{D}\Phi_1 D\Phi_1 + B\bar{D}C + 2\Psi_1^\dagger \bar{D}\Psi_1 + 2\Psi_2^\dagger \bar{D}\Psi_1 \right). \tag{3.7}$$

In terms of the free superfields the supercurrents can be written as

$$\begin{aligned}
\mathbf{J}^+ &= kDC + 2kCD\Phi_0 + 2C(\Psi_1\Psi_1^\dagger + \Psi_2\Psi_2^\dagger) + C^2B \\
&\quad - kD\Psi_1\Psi_1 - kD\Psi_2\Psi_2 - 2k\Psi_1\Psi_2D\Phi_1, \\
\mathbf{J}^- &= -B, \\
\mathbf{J}^0 &= BC + kD\Phi_0 + \Psi_1\Psi_1^\dagger + \Psi_2\Psi_2^\dagger, \\
\mathbf{J}^u &= -\Psi_1\Psi_2^\dagger + \Psi_2\Psi_1^\dagger - kD\Phi_1, \\
\mathbf{J}_1^{-\frac{1}{2}} &= \frac{1}{2}B\Psi_1 - \Psi_1^\dagger, \\
\mathbf{J}_2^{-\frac{1}{2}} &= \frac{1}{2}B\Psi_2 - \Psi_2^\dagger, \\
\mathbf{J}_1^{+\frac{1}{2}} &= kD\Psi_1 - k\Psi_1D\Phi_0 + C\mathbf{J}_1^{-\frac{1}{2}} - \Psi_1\Psi_2\Psi_2^\dagger + k\Psi_2D\Phi_1, \\
\mathbf{J}_2^{+\frac{1}{2}} &= kD\Psi_2 - k\Psi_2D\Phi_0 + C\mathbf{J}_2^{-\frac{1}{2}} + \Psi_1\Psi_2\Psi_1^\dagger - k\Psi_1D\Phi_1.
\end{aligned} \tag{3.8}$$

where

$$G_1 = \sin \Phi_1, \quad \lambda = \exp(\Phi_0).$$

3.2. HAMILTONIAN REDUCTION

Having constructed the phase space of the N=1 super Osp(2,2) in terms of free superfields, we now proceed to show that this phase space can be reduced to that of the N=2 superVirasoro algebra. The constraints we impose are the following,

$$\mathbf{J}^- = 0, \quad \mathbf{J}_1^{-\frac{1}{2}} = 0, \quad \mathbf{J}_2^{-\frac{1}{2}} = k. \quad (3.9)$$

An element of the space of the residual symmetry that preserves these constraints takes values in the Borel super subalgebra generated by $\{l^{-1}, l_1^{-\frac{1}{2}}, l_2^{-\frac{1}{2}}\}$ and is given by

$$U = \begin{pmatrix} 1 & 0 & 0 & 0 \\ A & 1 & \epsilon_1 & \epsilon_2 \\ \epsilon_1 & 0 & 1 & 0 \\ \epsilon_2 & 0 & 0 & 1 \end{pmatrix}. \quad (3.10)$$

Under the finite residual gauge transformation

$$g \rightarrow Ug, \quad (3.11)$$

the supercurrents transform as

$$\begin{aligned} \mathbf{J} \rightarrow & \begin{pmatrix} 1 & 0 & 0 & 0 \\ A & 1 & -\epsilon_1 & -\epsilon_2 \\ -\epsilon_1 & 0 & 1 & 0 \\ -\epsilon_2 & 0 & 0 & 1 \end{pmatrix} \mathbf{J} \begin{pmatrix} 1 & 0 & 0 & 0 \\ -A & 1 & -\epsilon_1 & -\epsilon_2 \\ -\epsilon_1 & 0 & 1 & 0 \\ -\epsilon_2 & 0 & 0 & 1 \end{pmatrix} \\ & + k \begin{pmatrix} 0 & 0 & 0 & 0 \\ DA - D\epsilon_2\epsilon_2 - D\epsilon_1\epsilon_1 & 0 & D\epsilon_1 & D\epsilon_2 \\ D\epsilon_1 & 0 & 0 & 0 \\ D\epsilon_2 & 0 & 0 & 0 \end{pmatrix}. \end{aligned} \quad (3.12)$$

If we select the gauge parameters as (Drinfeld-Sokolov gauge),

$$\epsilon_2 = \frac{1}{k}\mathbf{J}^0, \quad \epsilon_1 = -\frac{1}{k}\mathbf{J}^u, \quad A = -\frac{1}{k}\mathbf{J}_2^{+\frac{1}{2}} - \frac{1}{k}D\mathbf{J}^0, \quad (3.13)$$

then (3.12) can be used to bring $\mathbf{J}$ into the following form

$$\mathbf{J}_{DS} = \begin{pmatrix} 0 & 0 & 0 & k \\ X & 0 & Y & 0 \\ Y & 0 & 0 & 0 \\ 0 & -k & 0 & 0 \end{pmatrix}, \quad (3.14)$$

where X and Y , the coordinates of reduced phase space, are given by

$$\begin{aligned} X &= \mathbf{J}^+ + \frac{2}{k} \mathbf{J}_1^{+\frac{1}{2}} \mathbf{J}^u - \frac{2}{k} \mathbf{J}_2^{+\frac{1}{2}} \mathbf{J}^0 - D \mathbf{J}_2^{+\frac{1}{2}} - \partial_z \mathbf{J}^0 - \frac{1}{k} D \mathbf{J}^0 \mathbf{J}^0 - \frac{1}{k} D \mathbf{J}^u \mathbf{J}^u, \\ Y &= \mathbf{J}_1^{+\frac{1}{2}} - \frac{1}{k} \mathbf{J}^0 \mathbf{J}^u - D \mathbf{J}^u. \end{aligned} \quad (3.15)$$

Substituting the expressions of the supercurrents in (3.15) and using the constraints (3.9), we get

$$\begin{aligned} X &= -k(\partial_z \Phi_1 D \Phi_1 + \partial_z \Phi_0 D \Phi_0 + \partial_z D \Phi_0), \\ Y &= k(\partial_z \Phi_1 + D \Phi_0 D \Phi_1). \end{aligned} \quad (3.16)$$

In terms of the components,

$$\begin{aligned} \Phi_0 &= \phi_0 + \theta \psi_0, & \Phi_1 &= \phi_1 + \theta \psi_1, \\ X &= -(G_1 - \theta T), & Y &= U + \theta G_2, \end{aligned} \quad (3.17)$$

equation (3.16) gives,

$$\begin{aligned} T &= -k(\partial_z \psi_1 \psi_1 + \partial_z \psi_0 \psi_0 + \partial_z \phi_1 \partial_z \phi_1 + \partial_z \phi_0 \partial_z \phi_0 + \partial_z^2 \phi_0), \\ G_1 &= k(\partial_z \psi_0 + \psi_1 \partial_z \phi_1 + \psi_0 \partial_z \phi_0), \\ G_2 &= k(\partial_z \psi_1 - \psi_0 \partial_z \phi_1 + \psi_1 \partial_z \phi_0), \\ U &= k(\partial \phi_1 + \psi_0 \psi_1). \end{aligned} \quad (3.18)$$

Using

$$\begin{aligned} \{D \Phi_0(z_1, \theta_1), \Phi_0(z_2, \theta_2)\}_{\text{P.B}} &= -\frac{1}{2k} \theta_{12} \delta(z_{12}), \\ \{D \Phi_1(z_1, \theta_1), \Phi_1(z_2, \theta_2)\}_{\text{P.B}} &= -\frac{1}{2k} \theta_{12} \delta(z_{12}), \end{aligned} \quad (3.19)$$

the Poisson brackets of the coordinates $\{T, G_1, G_2, U\}$ can be easily shown to satisfy

the classical N=2 superconformal algebra,

$$\begin{aligned}
\{T(z_1), T(z_2)\}_{\text{P.B.}} &= \frac{k}{2} \partial_{z_2}^3 \delta(z_1 - z_2) + 2T(z_2) \partial_{z_2} \delta(z_1 - z_2) \\
&\quad + \partial_{z_2} T(z_2) \delta(z_1 - z_2), \\
\{T(z_1), G_1(z_2)\}_{\text{P.B.}} &= \delta(z_1 - z_2) \partial_{z_2} G_1(z_2) + \frac{3}{2} \partial_{z_2} \delta(z_1 - z_2) G_1(z_2), \\
\{T(z_1), G_2(z_2)\}_{\text{P.B.}} &= \delta(z_1 - z_2) \partial_{z_2} G_2(z_2) + \frac{3}{2} \partial_{z_2} \delta(z_1 - z_2) G_2(z_2), \\
\{T(z_1), U(z_2)\}_{\text{P.B.}} &= \delta(z_1 - z_2) U(z_2) + \partial_{z_2} \delta(z_1 - z_2) U(z_2), \\
\{G_1(z_1), G_1(z_2)\}_{\text{P.B.}} &= \frac{k}{2} \partial_{z_2}^2 \delta(z_1 - z_2) + \frac{1}{2} \delta(z_1 - z_2) T(z_2), \\
\{G_2(z_1), G_2(z_2)\}_{\text{P.B.}} &= \frac{k}{2} \partial_{z_2}^2 \delta(z_1 - z_2) + \frac{1}{2} \delta(z_1 - z_2) T(z_2), \\
\{U(z_1), G_1(z_2)\}_{\text{P.B.}} &= \frac{1}{2} \delta(z_1 - z_2) G_2(z_2), \\
\{U(z_1), G_2(z_2)\}_{\text{P.B.}} &= -\frac{1}{2} \delta(z_1 - z_2) G_1(z_2), \\
\{G_1(z_1), G_2(z_2)\}_{\text{P.B.}} &= -\frac{1}{2} \delta(z_1 - z_2) \partial_{z_2} U(z_2) - \partial_{z_2} \delta(z_1 - z_2) U(z_2), \\
\{U(z_1), U(z_2)\}_{\text{P.B.}} &= -\frac{k}{2} \partial_{z_2} \delta(z_1 - z_2).
\end{aligned} \tag{3.20}$$

The diagonal gauge is given by

$$\epsilon_1 = -\Psi_1, \quad \epsilon_2 = -\Psi_2, \quad A = -C. \tag{3.21}$$

In this gauge $\mathbf{J}$ takes the following form

$$\mathbf{J}_{\text{diagonal}} = \begin{pmatrix} kD\Phi_0 & 0 & 0 & k \\ 0 & -kD\Phi_0 & 0 & 0 \\ 0 & 0 & 0 & kD\Phi_1 \\ 0 & -k & -kD\Phi_1 & 0 \end{pmatrix}. \tag{3.22}$$

The coordinates $\{X, Y\}$ are connected to the coordinates $\{D\Phi_0, D\Phi_1\}$ by the super Miura transformation. This connection can be explained as in the previous cases

as follows. Consider the system of linear differential equation

$$(kD - \mathbf{J})v = 0, \quad v = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{pmatrix}. \quad (3.23)$$

Computing (3.23) in the diagonal gauge gives

$$\begin{aligned} kDv_1 - kD\Phi_0v_1 - kv_4 &= 0, \\ kDv_2 + kD\Phi_0v_2 &= 0, \\ kDv_3 - kD\Phi_1v_4 &= 0, \\ kDv_4 + kv_2 + kD\Phi_1v_3 &= 0. \end{aligned} \quad (3.24)$$

By solving these equation in terms of v_1 , the following gauge-invariant differential equation is obtained,

$$k\left(\partial_z D - (\partial_z \Phi_0 v_1 + D\Phi_0 \partial_z \Phi_0) + k(\partial_z \Phi_1 + D\Phi_0 D\Phi_1) \frac{1}{D}(\partial_z \Phi_1 + D\Phi_0 D\Phi_1)\right)v_1 = 0. \quad (3.25)$$

In the Drinfeld-Sokolov gauge, (3.23) becomes

$$\begin{aligned} kDv_1 - kv_4 &= 0, \\ kDv_2 - Xv_1 - Yv_3 &= 0, \\ kDv_3 - Yv_1 &= 0, \\ kDv_4 + kv_2 &= 0. \end{aligned} \quad (3.26)$$

Similarly, by solving these equations in terms of v_1 we get

$$k\partial_z Dv_1 + Xv_1 + Y\frac{1}{D}(Yv_1) = 0. \quad (3.27)$$

Comparing (3.25) and (3.27) we get the expressions of X and Y in (3.16).

4. Discussions

In this letter, the Hamiltonian reduction of the $SL(2, \mathbb{R})$ current algebra is reviewed using its free-field realization. We also have demonstrated the equivalence of this formalism to Polyakov “Soldering procedure”. The results of [8] were also verified using the free-superfield realization of the $N=1$ $Osp(1,2)$ supercurrent algebra. Motivated by the results of [9] in which the Polyakov method has been employed to extract the $N=2$ superconformal transformations from the $N=1$ $Osp(2,2)$ supergauge transformation, we have considered the classical Hamiltonian reduction of the $N=1$ $Osp(2,2)$ supercurrent algebra using its free-superfield realization. The reduced phase space was shown to exhibit the classical $N=2$ superconformal algebra. Our construction can be extended to the cases of $N=1$ $Osp(n, 2)$ ($n > 2$) supercurrent algebras. We expect that these algebras will reduce to the $O(n)$ -extended superconformal algebras found by Bershadsky and Knizhnik [19].

Finally we give some comments on the quantum theory. The quantum WZNW model with a group G is a superconformal and super G Kac-Moody invariant two dimensional field theory. The super stress-energy tensor $\mathcal{T}$ constructed from the super Kac-Moody currents via the super Sugawara construction [17,18] together with the supercurrents satisfy the following operator product expansion

$$\begin{aligned}\mathcal{T}(z_1, \theta_1)\mathcal{T}(z_2, \theta_2) &\sim \frac{c}{6} \frac{1}{z_{12}^3} + \frac{3}{2} \frac{\theta_{12}}{z_{12}^2} \mathcal{T}(z_2, \theta_2) + \frac{1}{2z_{12}} D\mathcal{T}(z_2, \theta_2) + \frac{\theta_{12}}{z_{12}} \partial\mathcal{T}(z_2, \theta_2), \\ \mathcal{T}(z_1, \theta_1)\mathbf{J}^{(a)}(z_2, \theta_2) &\sim \frac{1}{2} \frac{\theta_{12}}{z_{12}^2} \mathbf{J}^a(z_2, \theta_2) + \frac{1}{2} \frac{D_2 \mathbf{J}^a(z_2, \theta_2)}{z_{12}} + \frac{\theta_{12} \partial_{z_2} \mathbf{J}^a(z_2, \theta_2)}{z_{12}}, \\ \mathbf{J}^a(z_1, \theta_1)\mathbf{J}^b(z_2, \theta_2) &\sim \frac{k/2\eta^{ab}}{z_{12}} + \frac{\theta_{12} f^{ab}{}_c \mathbf{J}^c(z_2, \theta_2)}{z_{12}},\end{aligned}\tag{4.1}$$

where η^{ab} and $f^{ab}{}_c$ are the metric and structure constant of the Lie algebra $\mathcal{G}$ of G . The central charge of the superVirasor algebra is given by

$$c = \frac{3}{2} \left(1 - \frac{2C_{\text{adj}}}{k} \right) \dim \mathcal{G},\tag{4.2}$$

where C_{adj} is the dual Coxeter number of the group G and $\dim$ is the dimension

of the group G (in the case when G is a supergroup the dim is replaced by the $sdim$, where $sdim$ is the dim of the bosonic generators minus the dim of the fermionic generators of the group G). The second equation in (4.1) means that all the supercurrents have the dimension $1/2$ with respect to the super stress-energy tensor $\mathcal{T}$.

In the analysis of [4] where the quantum Hamiltonian reduction of the $SL(2, \mathbb{R})$ bosonic current algebra is considered, the stress tensor has to be modified in order that the constraints are consistent with conformal invariance. However, in the case of the Hamiltonian reduction of the bosonic $Osp(1,2)$ current algebra [5], after imposing the constraint $J^- = 1$ and modifying the stress tensor as in the case of $SL(2, \mathbb{R})$, additional Majorana fermion has to be introduced for the closure of the algebra of constraints. Similarly, in the Hamiltonian reduction of $Osp(2,2)$ current algebra two Majorana fermions have to be introduced in order that the constraints form a closed algebra.

In [15] where the quantum Hamiltonian reduction of the $Osp(1,2)$ supercurrent algebra is considered, an appropriate super stress-energy tensor is constructed by modifying the Sugawara form of the super stress-energy tensor as

$$\mathcal{T}' = \mathcal{T} - \partial_z \mathbf{J}^0. \quad (4.3)$$

In this construction no additional structure has to be introduced as the superspace constraints are consistent and form a closed algebra. This is because the superspace current $\mathbf{J}^{(-\frac{1}{2})}$ is of dimension 0 with respect to $\mathcal{T}'$.

The same situation obviously arises in the quantum Hamiltonian reduction of $Osp(2,2)$ supercurrent algebra. After modifying the super stress-energy tensor as in (4.3), the supercurrents $\mathbf{J}_1^{(-\frac{1}{2})}$ and $\mathbf{J}_2^{(-\frac{1}{2})}$ are both of dimension 0 and the algebra of constraints closes without introducing extra fermions. Then following the standard procedure of quantizing a system with constraints, one introduces three superspace ghost systems (corresponding to the three constraints), $(b(z, \theta), c(z, \theta))$,

$(\eta(z, \theta), \zeta(z, \theta))$ and $(\eta'(z, \theta), \zeta'(z, \theta))$ with dimensions $(-\frac{1}{2}, 1)$, $(0, \frac{1}{2})$, and $(0, \frac{1}{2})$ respectively. In components, these superghosts are given as

$$\begin{aligned} b(z, \theta) &= b_1 + \theta b_2, & c(z, \theta) &= c_1 + \theta c_2, \\ \eta(z, \theta) &= \eta_1 + \theta \eta_2, & \zeta(z, \theta) &= \zeta_1 + \theta \zeta_2, \\ \eta'(z, \theta) &= \eta'_1 + \theta \eta'_2, & \zeta'(z, \theta) &= \zeta'_1 + \theta \zeta'_2. \end{aligned} \tag{4.4}$$

The ghosts (b_2, c_1) , (η_1, ζ_2) and (η'_1, ζ'_2) are all anticommuting and have the dimension $(0, 1)$ while the pairs (b_1, c_2) , (η_2, ζ_1) and (η'_2, ζ'_1) are all commuting and have the dimensions of $(-\frac{1}{2}, \frac{3}{2})$, $(\frac{1}{2}, \frac{1}{2})$ and $(\frac{1}{2}, \frac{1}{2})$ respectively. The total super stress-energy tensor $\mathbf{T}^{\text{total}}$ of the system is the sum of the modified super stress energy tensor and that of the superghosts. The superghosts contribute +3 to the superconformal anomaly. Thus the total central charge of the reduced theory is given by

$$c^{\text{tot}} = -6k + 3. \tag{4.5}$$

In order to complete the quantum Hamiltonian reduction, the reduced Hilbert space must be obtained. One has to construct the BRST operator and study its cohomology following [4,5,15]. Then this cohomology must be shown to be isomorphic to an irreducible representation space of the N=2 superVirasoro algebra with central charge $c = 3p/(p+2)$ (where $k = 1/(p+2)$). We hope to report on this in a future publication.

ACKNOWLEDGEMENT

I would like to thank T. Kuramoto for useful discussions.

REFERENCES

1. A. A. Belavin, in *Quantum String Theory*, Volume 31, Springer-Verlag, Berlin, Heidelberg, 1988 page 132.
2. A. Alekseev and Shatashvili, Nucl. Phys B323 (1989) 719.
3. V. Drinfeld and V. Sokolov, Journal of Soviet Mathematics 30, (1984) 1975.
4. M. Bershadsky and H. Ooguri, Comm. Math. Phys. 2 (1989) 49.
5. M. Bershadsky and H. Ooguri, Phys. Lett, 229B (1989) 374.
6. A. M. Polyakov, Int. J. Mod. Phys. A5(1990) 833.
7. A.M. Polyakov, Mod. Phys. Lett. A2 (1987) 893;
V.G. Knizhnik, A.M. Polyakov and A.B. Zamolodchikov, Mod. Phys. Lett. A3 (1988) 819;
A.M. Polyakov and A.B. Zamolodchikov, Mod. Phys. Lett. A3 (1988)1213;
Lectures given by Polyakov at Les Houches summer school on Fields, Strings and Critical Phenomena (1988).
8. W. A. Sabra, Nucl. Phys B375 (1992) 82.
9. W. A. Sabra, Restricted gauge invariance, N=2 Coadjoint orbits and N=2 quantum supergravity, preprint, BIR/PH/92-1.
10. M. Wakimoto, Comm. Math. Phys. 104 (1986) 605.
11. A.B. Zamolodchikov, unpublished
12. S. P. Novikov, Sov. Math. Dokl. 24 (1981) 222.
13. E. Witten, Comm. Math. Phys. 92 (1984) 455.
14. A.M. Polyakov and P.B. Wiegmann, Phys. Lett. 141B (1984) 233.
15. T. Kuramoto, Quantum Hamiltonian Reduction Of Super Kac-Moody Algebra II, preprint, YNUE-PH-92-01.
16. B. L. Feigin and D. B. Fuchs, Funct. Anal. and Appl. 16 (1982) 114; 17 (1983) 241.

17. P. di Vecchia, V. G. Knizhnik, J. L. Peterson and P. Rossi, Nucl. Phys B253 (1985) 701.
18. J. Fuchs, Nucl. Phys B286 (1986) 455, B318 (1989) 631.
19. V. G. Knizhnik, Theor. Math. Phys. 66 (1986) 68; M. Bershadsky, Phys. Lett. 174B (1986) 285.